## 耳介下部膿瘍摘出術後の 下顎・側頸部の難治性疼痛の1治験例

## 林 明宗\*

要旨: 左耳介下部膿瘍切除術後に続発した,左下顎・側頸部・前胸部にわたる激しい痛みを訴えた患者の1治験例を報告する.星状神経節ブロックが有用であったが,満足できるものではなかった. 当帰四逆加呉茱萸生姜湯(TJ-38),桂枝茯苓丸(TJ-25)などの駆芎血剤と筋弛緩剤との併用が,満足できる除痛をもたらすことができた.

索引用語:頑痛. 当帰四逆加呉茱萸生姜湯. 桂枝茯苓丸

## PAIN AND KAMPO MEDICINE Vol. 13 (2003)

A case of intractable pain in the submandibular and lateral cervical region after infraauricular abscess resection

Akimune Hayashi\*

**Abstract**: The author report on a patient suffering from intractable pain in the submandibular, lateral cervical and anterior thoracic region after surgical extirpation of the left infraauricular abscess. Satellite ganglion block was somewhat effective but not satisfactory. The following Kampo prescriptions combined with a muscle relaxant were successful in resolving the pain satisfactorily: Toki-shigyaku-ka-gosyuyu-syokyo-to (TJ-38) and Keishi-bukuryo-gan (TJ-25).

**Key words**: Intractable pain, Toki-shigyaku-ka-gosyuyu-syokyo-to (TJ-38), Keishi-bukuryo-gan (TJ-25)

<sup>\*</sup>Department of Neurosurgery, Yokohama City University Medical Center, Yokohama City University School of Medicine

Offprint requests to: Akimune Hayashi, Department of Neurosurgery, Yokohama City University Medical Center, Yokohama-City University School of Medicine, 4-57, Urafune-Cho, Minami-Ku, Yokohama, 233-0024, Japan