臍帯血移植後に発症した 左側腹部〜腰背部頑痛に対する漢方治療

林 明宗*

要旨:近代西洋医学の抗癌治療は、多くの合併症を引き起こす。今回、抗癌治療の後に著明な腎虚による頑痛症状を呈し、補腎治療が奏功した1例を経験したので報告する。症例は28歳男性で、急性リンパ球性白血病に罹患している。臍帯血輸血を施行されたが、治療後抗癌剤と免疫抑制剤の服用を余儀なくされた。このため極度の腎気虚症状による腰背部頑痛と合併する脾虚症状を呈してきた。これに対して牛車腎気丸と修治附子末を投与したところ、80%以上の症状改善が認められた。

索引用語:急性リンパ球性白血病, 頑痛, 牛車腎気丸, 修治附子末N, 十全大補湯

PAIN AND KAMPO MEDICINE Vol. 16 (2006)

Oriental Medical Therapy for Intractable Lateral Abdominal and Low Back Pain after Umbilical Cord Blood Transplantation Akimune Hayashi*

Abstract: Modern western anti-cancer therapy can cause several complications. We experienced a case of severe kidney-Ki deficiency symptoms after anti-cancer therapy. The case showed excellent response to nourishing kidney-ki prescriptions. A 28-year-old male patient suffering from acute lymphocytic leukemia underwent umbilical cord blood transplantation and had to take anti-cancer and immunosuppressant therapy thereafter. He complained of intractable low-back pain due to severe kidney-ki deficiency combined with spleen-ki deficiency symptoms. Goshajinkigan and shujibushimatsu N had resolved his symptoms more than 80%.

Key words: acute lymphocytic leukemia, intractable pain, goshajinkigan, shujibushimatsu N, juzentaihoto

^{*} Department of Neurosurgery, Kanagawa Cancer Center, Yokohama, Japan Division of Kampo Medicine, Department of General Internal Medicine, Yokohama City University Medical Center

Offprint requests to: Akimune Hayashi, Department of Neurosurgery, Kanagawa Cancer Center, Yokohama, Japan. 1-1-2 Nakao, Asahi-Ku, Yokohama, 241-0815, Japan