臨床経験

胆嚢癌肝拡大右葉切除術後の上腹部痛に対し 芍薬甘草湯が有効であった一例

円城寺昭人*1 川口真規子*1 今村一歩*1 中島弘治*1 山口 泉*1 森内博紀*1 山口淳三*1 黒木 保*2 田島義証*2 兼松隆之*2 佐藤英俊*3

要旨: 症例は 68 歳女性, 胆嚢癌肝浸潤の診断にて肝拡大右葉切除術を施行後, 腹腔動脈周囲リンパ節再発, 残肝再発した. 術後 3 週間目より右季肋部鈍痛と右側腹部にいたる筋痙攣性の痛みに対し, 抗コリン性鎮痙薬, NSAIDs を使用したが難治であった. 術後早期には画像上再発を確認できなかったため, オピオイドの使用が遅れたが芍薬甘草湯(TJ-68) を投与したところ有効であった. がん性疼痛では痛みの性質がさまざまで筋痙攣性の疼痛には芍薬甘草湯などの漢方が奏効する可能性がある.

索引用語:胆囊癌再発、癌性疼痛、筋痙攣性、芍薬甘草湯

PAIN AND KAMPO MEDICINE Vol.20 (2010)

A case report of shakuyakukanzoto which can control the severe abdominal pain caused by the recurrence of gallbladder carcinoma after the radical resection Akihito Enjoji *1, Makiko Kawaguchi *1, Hajime Imamura *1, Hiroharu Nakajima *1, Izumi Yamaguchi *1, Hironori Moriuchi *1, Junzo Yamaguchi *1, Tamotsu Kuroki *2, Yoshitsugu Tajima *2, Takashi Kanematsu *2 and Hidetoshi Sato *3

Abstract: A case was the female patient who had the recurrence of gallbladder carcinoma after the radical resection. She complained of severe abdominal pain come from the recurrence site at lymph nodes around celiac ganglion. Although the pain was uncontrollable by opioid and various analgesic agents, shakuyakukanzoto (TJ-68) could relieve the abdominal pain caused by the recurrence of gallbladder carcinoma after the radical resection.

Key words: shakuyakukanzoto, recurrence of gallbladder carcinoma, cancer pain

^{* 1} National Hospital Organization Saga National hospital Offprint requests to: Akihito Enjoji, National Hospital Organization Saga National hospital. 1-20-1, Hinode, Saga, 849-0923, Japan

^{* &}lt;sup>2</sup> Department of Surgery, Nagasaki University Graduate School of Biomedical Sciences

^{* &}lt;sup>3</sup> Department of Community-based Comprehensive Palliative, Care Saga Medical School Hospital, Saga University