症例報告

帯状疱疹痛に対する柴胡疎肝湯の効果 - 3 症例の検討-

要旨: 帯状疱疹痛に対する柴胡疎肝湯の効果について検討した. 帯状疱疹発症後 6-23 日の亜急性の帯状疱疹痛を有する3症例に対して柴胡疎肝湯をエキス剤の合剤で投与し, 奏効した. 帯状疱疹に関する痛みは治療が困難なことがある. 帯状疱疹後神経痛への移行 を阻止するためにも初期の鎮痛法の一つとして柴胡疎肝湯は有効である.

索引用語:柴胡疎肝湯,带状疱疹後疼痛,带状疱疹後神経痛

PAIN AND KAMPO MEDICINE Vol.21 (2011)

Effect of saikosokanto on the post hepatic pain in 3 cases.

Satoko Uemura *1, Ryo Katsuki *2, Naomi Hirakawa *1 and Michihiko Hirata *3

Abstract: Post hepatic pain (PHP) is a difficult pain to control. Administration of saikosokanto was effective for the severe PHP in 3 patients. To treat the PHP is important to prevent the post hepatic neuralgia. It was suggested that the administration of saikosokanto should be the good therapy to improve PHP.

Key words: saikosokanto, Posthepatic Pain, Posthepatic Neuralgia

^{* 1} Department of Anesthesiology, Saga Medical School Offprint requests to: Satoko Uemura, Department of Anesthesiology, Saga Medical School. Nabeshima 5-1-1, Saga, 849-8501, Japan

^{* &}lt;sup>2</sup> Department of Anesthesia, NHO Ureshino Medical Center

^{* 3} Hirata Pain Clinic