

症 例 報 告

難治性吃逆に対し半夏瀉心湯が有効であった 1 症例

石川亜佐子*¹ 香月 亮*¹ 平田道彦*²

要旨：肝癌の治療中，肝腫瘍による難治性吃逆を発症した症例を経験した．芍薬甘草湯を投与したが効果がなく，肝腫瘍による上腹部の鞭満所見を心下痞鞭と捉え，半夏瀉心湯を選択し，有効であった．心下痞鞭のある吃逆患者については，半夏瀉心湯が有効な場合がある．

索引用語：吃逆，半夏瀉心湯，心下痞鞭，肝腫瘍

PAIN AND KAMPO MEDICINE Vol.24 (2014)

A case of refractory hiccough treated with hangeshashinto

Asako ISHIKAWA*¹, Ryo KATSUKI*¹ and Michihiko HIRATA*²

Abstract: We reported a case of refractory hiccough caused by liver cancer under the medical treatment. Since shakuyakukanzoto had no effect on the hiccough, hangeshashinto was administered from a viewpoint of the fullness of abdominal area according to the Kampo theory. The hiccough was gradually disappeared from the second medication. It was suggested that the fullness of the upper abdominal area should be the important sign at the time of choosing hangeshashinto.

Key words: hiccough, hangeshashinto, shinkahiko, liver cancer

*¹ NHO Ureshino Medical Center

Offprint requests to: Asako Ishikawa, NHO Ureshino Medical Center,
2436 Hei Oazashimajuku, Ureshino City, Saga 843-0393, Japan

*² Hirata Painclinic