症例報告

膵尾部粘液性嚢胞性腺腫摘出術後の遷延性背部痛に 漢方治療が有用であった1症例

林 明宗*

要旨:55歳女性. 膵尾部粘液性嚢胞性腺腫摘出術の術後に仮性嚢胞が形成されるとともに, 背部痛を合併し, 疼痛は4年余にわたって遷延した. 術後の組織癒着, 仮性嚢胞による周辺組織の圧迫によって生じた局所の瘀血, 浮腫ならびに気虚に対する漢方治療を行い, 症状の軽快を認めた.

索引用語: 膵嚢胞, 術後疼痛, 漢方治療

PAIN AND KAMPO MEDICINE Vol.25 (2015)

Kampo therapy for prolonged intractable low-back pain after resection of mucinous cystadenoma from the pancreatic tail

Akimune HAYASHI*

Abstract: A 55-year-old woman underwent an operation for mucinous cystadenoma in the pancreatic tail. Due to a complication from the surgery, she developed postoperative pancreatic pseudocyst and had been suffering from prolonged low-back pain for more than four years. Kamopo therapy treated local blood stasis, edema (fluid retention), and qi deficiency, which were caused by postoperative tissue adhesion and scar formation, and the compression of the surrounding tissue by pancreatic pseudocyst. The treatment successfully relieved the patient's complaints.

Key words: pancreatic cyst, post-operative pain, Kampo therapy

^{*} Department of Oriental Medicine, Kampo Support Center, Kanagawa Cancer Center

Offprint requests to: Akimune Hayashi, Department of Oriental Medicine, Kampo Support Center, Kanagawa
Cancer Center.

²⁻³⁻² Nakao, Asahi-Ku, Yokohama, Kanagawa 241-8515, Japan